




Do I need to tell my sexual partners that I have been diagnosed with chlamydia?

Yes, you will need to tell any sexual partners from the last six months so that they can be tested and treated. Telling sexual partners is the best way to control the spread of chlamydia.

How do I tell them?

It is best to tell partners directly, either in person, by phone call, or text message. Your nurse or doctor can help you with this if needed.

There are also websites which allow you to anonymously notify someone that they have been exposed to an STI. These websites are:



Let them Know
letthemknow.org.au
This website is if available
for anyone to use.



The Drama Down Under
This website is specifically
for gay men, or men with
male sexual partners,
and allows you to contact
several people in one go.

Where to go for information and testing:

- SEXUAL HEALTH & FAMILY PLANNING ACT (SHFPACT)
Call 026247 3077 or visit shfpact.org.au
- Canberra Sexual Health Centre (Canberra Hospital)
Call 02 5124 2184 or visit health.act.gov.au/sexualhealth
- ACT Walk-in Centres (can provide testing for chlamydia)
- Your own GP or local health centre



SHFPACT Clinical Services

- Contraceptive Advice, Information & Services
- IUD's and other Long Acting Contraceptive Options
- Cervical Screening
- Canberra Menopause Centre
- Pregnancy Options Counselling
- Sexually Transmissible Infections(STIs):
- Testing, Treatment and Advice
- Blood borne virus and HIV testing
- Emergency Contraception (the 'Morning After' Pill)
- Breast Checks and Breast Awareness Education
- Fertility Issues and Pre-conception Advice
- Sexual Difficulties and Sexuality Education
- PrEP Information, Advice, and Prescription

References: Australian STI Management Guidelines for use in primary care.

<https://sti.guidelines.org.au/>
Stay STI Free-Victorian Sexual Health Network:
<https://www.staystifree.org.au/get-the-facts/chlamydia>




August 2015

SHFPACT is a health promotion charity (HPC) assisted with funding from the ACT Government. All donations to SHFPACT over \$2.00 are tax deductible.



Chlamydia



Chlamydia is the most common bacterial STI in young people under 30 in Australia. It is easy to catch, easy to test, and easy to treat.

Level 1, 28 University Ave
Canberra ACT 2601
shfpact@shfpact.org.au
02 6247 3077
shfpact.org.au





What is chlamydia?

Chlamydia is a very common sexually transmissible infection (STI) which is caused by a bacterium called chlamydia trachomatis.

It is the most common bacterial STI in young people under 30 in Australia.

Chlamydia is easy to catch, easy to test, and easy to treat.

It can cause infection of the cervix, anus, throat, urethra (penis), and occasionally the eyes.

How do you get chlamydia?

You get chlamydia by having sex without a condom with a person who has the infection. Chlamydia is sometimes also transmitted through oral sex.

Is a chlamydia infection serious?

If chlamydia is left undetected and untreated, it can cause serious health problems; this includes Pelvic Inflammatory Disease (PID). PID is a serious infection of the uterus (womb) and the fallopian tubes and can lead to chronic pelvic pain and infertility.

Chlamydia infection can also cause damage to the small tubes in the testes which can also affect fertility.

Early detection and treatment can prevent these problems occurring.

What are the symptoms of chlamydia?

Most people with chlamydia have no symptoms. This is why it's important to test regularly. If symptoms do occur, they include:

If you have a vagina:

- Unusual vaginal bleeding (between periods or after sex)
- Unusual vaginal discharge
- Pelvic pain
- Pain during sex
- Pain or burning passing urine

If you have a penis:

- A discharge from the penis (often clear)
- Redness at the tip of the penis
- Pain and swelling in the testicles
- Pain or burning passing urine

Anus:

- Pain or bleeding from the anus

How do you test for chlamydia?

Testing for chlamydia is easy.

- If you have a penis, the test for chlamydia is a urine test.
- If you have a vagina, the test for chlamydia is a self-collected vaginal swab or a urine test.
- If you are having anal sex, you may also need a self-collected anal swab.

When should I have a test for chlamydia?

You should have a chlamydia test if:

- You have ever had unprotected sex
- You have had a new sexual partner since your last chlamydia test
- You have a sexual partner who has been diagnosed with chlamydia or another STI
- Your sexual partner has had sex with a person who may have chlamydia
- You have been diagnosed with another STI
- You have any signs or symptoms

How do you treat chlamydia?

Chlamydia is treated with oral antibiotics. If you have symptoms, you may need different antibiotics, your doctor will discuss this with you.

How long does it take after treatment for any symptoms to go away?

If you have symptoms, they should start to go away within a few days of starting treatment.

When can I have sex again after treatment for chlamydia?

It is essential that you do not have any sex until 7 days after you finish your treatment.

If your partner has been treated, you must not have sex with them again until 7 days after they finish their treatment.

Do I need to be tested again?

Yes, you should have a test three months after treatment. This is to make sure that you have not been re-infected.

How do I protect myself from chlamydia?

The best way to protect yourself from chlamydia infection is to use condoms when having sex (and a dam for oral sex).

To fully protect yourself, you need to use a condom the whole time you have sex, so you avoid sharing any body fluids.

Sex toys also need to be washed or use a condom to cover them before sharing between partners.

