

# Pregnancy Options

for an unintended pregnancy

Level 1, 28 University Ave  
Canberra ACT 2601  
[shfpact@shfpact.org.au](mailto:shfpact@shfpact.org.au)  
02 6247 3077  
[shfpact.org.au](http://shfpact.org.au)





Unintended pregnancy is common. It is estimated that 150,000 to 200,000 unintended pregnancies occur in Australia every year.

There are three choices available when you have an unintended pregnancy:

- Continue the pregnancy and parent
- Continue the pregnancy and place the child for adoption or kinship care
- Have an abortion

### What to do first if you think you might be pregnant

If you think you are pregnant the first step is to confirm if you are. To do this you need to have a pregnancy test. Confirming if you are pregnant is important so that you can get the right care early if you are continuing the pregnancy, access timely abortion if you are not continuing, and give yourself time to decide if you are unsure.

a urine pregnancy test after you have missed your period, or 16 days after the sex you are worried about. If you do a test too early, such as before a missed period, it may not be correct.

If the pregnancy test is positive, you can confirm the pregnancy with your GP, or a nurse or doctor at a family planning clinic. Seeing a nurse or doctor will allow you to obtain further information, advice, assistance, or referral as needed. They can also do a pregnancy blood test if needed, which is more accurate than a urine test and can give an estimation of how far pregnant you are.

If you have symptoms of pregnancy, have had at least one positive home pregnancy test, and know that you do not want to continue the pregnancy, you do not need to see a GP, you can book into an abortion clinic without a referral.

If the pregnancy test is negative and your period still has not come, repeat the test again in a week. If it is still negative, then you should visit your GP or a family planning clinic to discuss this.

### Where can I have a pregnancy test?

You can do a home urine pregnancy test, or you can go to your GP or a family planning clinic to have a test. Home pregnancy tests are available in supermarkets and pharmacies. If you make sure that the test is in date and follow the manufacturer's instructions correctly, the accuracy of these tests is around 97%. Using the first urine sample in the morning can make it more accurate. Remember that you need to do

## What are the symptoms of pregnancy?


The symptoms of pregnancy can vary. The most common symptom that occurs is no period, or your period being unusually short and light.

### Other symptoms include:

- Swollen and tender breasts
- Nipple sensitivity
- Tiredness
- Urinating more often
- Nausea
- Vomiting
- Food aversions (finding the taste and smell of some foods unpleasant)
- Food cravings
- Mild pelvic cramping which may feel like period cramps
- Feeling more emotional or moody

Some people will get just a few symptoms, some will get many or all these symptoms, and for others the only symptom they may have is missing a period.

referred to as 'gestation'. For example: 8 weeks and 4 days gestation.



### Pregnancy is divided into 3 trimesters:

- The **first trimester** is from the beginning to 12 weeks.
- The **second trimester** is from 13 weeks to 28 weeks.
- The **third trimester** is from 29 weeks to 40 weeks.

## Dealing with an unintended pregnancy

Experiencing an unintended pregnancy can be a stressful and confusing time. Shock is a common experience when you first find out that you are pregnant, even if you already suspected that you were. You might find that you cannot think clearly and may experience a lot of mixed emotions. This feeling of shock may continue for some days.

Feelings of uncertainty and ambivalence are also very common, and this can occur even when a pregnancy is planned.

An unintended pregnancy can be particularly difficult if you are feeling uncertain about what decision you will make about the pregnancy, or if your thoughts and feelings about what you want to do differ from that of your partner, or the conception partner.

Emotions can be quite intense during pregnancy. This is thought to be because of pregnancy hormones. You may experience mood swings and may become upset more easily. This can make dealing with an unintended pregnancy and making an important decision like this more challenging.

## How will I know how pregnant I am?

A pregnancy is calculated from the first day of your last normal menstrual period. If you see a doctor or nurse, they can help you to work out how pregnant you are using this date.

They may also use the results of a pregnancy blood test to work it out, and possibly an ultrasound if you have an irregular menstrual cycle or don't know when your last period was.

If you are going to have an ultrasound and you are unsure if you are continuing the pregnancy, tell the doctor and the technician doing the ultrasound so that they can approach the ultrasound sensitively and appropriately.

Pregnancy is counted in weeks and days rather than months, and the length is

## Managing stress

The following are some ways that can help you to manage stress. These are good techniques to use in normal daily life and are particularly useful during a difficult time when you are feeling stressed or upset:

### Dropping anchor

Sitting quietly, first notice the thoughts and feelings that you are having and acknowledge that they are there. Then bring your awareness back into your body, straighten your back, push your feet into the floor, push your fingertips together, move your shoulders, elbows, and arms. Then look around you, notice what you can see, what you can hear, what you can smell, what you can feel, and what you can taste. Now notice your breathing, and slow and soften it. This exercise can help to ground you and bring you back into the present when you are feeling stressed and caught up in difficult thoughts or feelings.

### Slow breathing

Using your abdominal muscles inhale to the count of 4, pause and then exhale to the count of 8, repeat. Try to do this for a few minutes and do it regularly throughout the day or when you are feeling stressed.

### Box breathing

Inhale to the count of 4, pause to the count of 4, exhale to the count of 4, pause to the count of 4, repeat. Do this exercise for at least a few minutes.

### Make room for the feelings

Make time to sit down somewhere quiet and focus on the feelings that you are experiencing. Focus on one feeling at a time. Be curious about that feeling, where is it in your body? Is it big or small? Does it have firm edges or soft edges? What shape is it? Does it have a colour? Breathe into the feeling, soften around it, and make room for it to be there. This can help you to be more comfortable with these feelings and allow you to think more clearly, and to then take the action that you need to.

## Dealing with thoughts

Our minds are very busy, and at times of stress this is particularly so. This is very normal. Learning to notice your thoughts can help you to not become hooked by them or overwhelmed by them. When having a thought, ask yourself if it is a helpful or unhelpful thought. If it is helpful you may want to pay attention to it, if it is unhelpful or is not a good time to have this thought, then you can use some techniques to lessen the effect of that thought. You can notice a thought and say to yourself 'I notice I'm having the thought that...' then bring yourself back into the present moment. Then this just becomes another thought. You can also just notice thoughts like leaves floating past on a stream, or clouds drifting past in the sky, and again bring yourself back into the present.

### Self-care

It's very important to remember to look after yourself during a difficult and stressful time.

- Acknowledge that this is a stressful time for you
- Take time out from focussing on the decision
- Nurture yourself by taking time to relax, to rest and to do some of the things that you enjoy and that nourish you, such as taking a bath, going for a long walk, having a massage or a facial, going to a movie, or catching up with close friends.
- Practising self-compassion, meaning kindness towards yourself, is also very important at a time like this.



## Getting support

Talking to someone close who you trust and who you know will support you in whatever decision you make can also be very helpful.

If you are feeling particularly distressed and don't have anyone that you can talk to it's important that you seek help: see your GP; make an appointment with a known psychologist if you have one; make an appointment to see a pregnancy options counsellor (see details below); or call **Lifeline** on **13 11 14**.

You can also access **Medicare Mental Health**, a free service providing support for people with mental health concerns and connecting them to the right service.



If you are concerned about domestic violence and staying safe, call the National Domestic Family and Sexual Violence Counselling Service at **1800 RESPECT (1800 737 732)** or the **Domestic Violence Crisis Service** on **6280 0900** if you are in the ACT.

**If you are in immediate danger, call 000.**



## Pregnancy options and making a decision

For some people having to decide whether to continue a pregnancy or not can be relatively straightforward. For others it may be a difficult decision depending on their individual circumstances.

When considering your options in relation to a pregnancy you may consider many factors, including:

- The state and stability of your relationship with the conception partner
- The level of support you may have from family or friends
- Whether you feel ready to take on a parenting role
- The needs of any children you may already have
- Your career or study plans and goals
- Your housing and financial situation
- Your life plans and goals
- Your physical and emotional health
- Your beliefs and values

Unless you feel quite clear about what you are going to do it is usually advisable to take time to make a decision. Remember however that this is a time sensitive decision, and you may need to decide relatively quickly if your pregnancy is more advanced.

Consider the different options and factors impacting your decision, and if you are in a relationship talk it through with your partner.

If you find that it is just too difficult and you are unable to decide, or you would like to talk it through with somebody to clarify your thoughts and feelings, counselling may be helpful for you.

When you have decided about a pregnancy, whatever that decision is, make sure you write down the reasons for your decision and keep it in a safe place. If you forget the reasons why you made a certain decision in the future, and question it, this will help remind you why this was the right decision for you at the time.



### Deciding as a couple

While the decision whether to continue a pregnancy is essentially your decision as the pregnant person, if you are in a relationship, it is a decision that will inevitably affect you as a couple.

Talking about an unintended pregnancy as a couple can be difficult. Before you do this, you might want to think about your own feelings about the pregnancy first, so you have some clarity before you involve your partner. It may also be helpful to think about what you want from your partner before talking to them. You might want to involve them in the decision, just tell them about the decision that you have made, or just make time to listen to each other's views even if you feel they will not support your decision.

When you do talk about it as a couple, make sure that you allow focused time to do this, and agree to take turns to fully and respectfully listen to each other so that you both feel heard.

### For further information about making a decision



**Children by Choice**  
<https://www.childrenbychoice.org.au/information-support/decision-making/>

## Continuing the pregnancy and parenting

Choosing to become a parent, or to raise another child, is a life-changing decision. About half of those who experience an unintended pregnancy choose to continue the pregnancy and parent. There are many issues that you may consider when thinking about this option, these may include:

- Your existing relationship and individual family circumstances
- The level of support from your partner
- The level of parenting support you may get from the conception partner if they are not your partner
- How you feel about a continuing connection with the conception partner if they are not your partner
- Support that you may have from your own or your partner's parents, extended family, or friends
- Your financial situation
- Whether you have secure housing
- Study, employment, and career issues
- The use of drugs and alcohol during pregnancy
- How you would look after a baby & child - feeding, health, and wellbeing
- Any domestic violence and issues relating to your own and any child's safety and wellbeing

### Questions you may want to ask yourself if you are considering continuing the pregnancy and parenting

- How do I feel about the prospect of becoming a parent/or having another child?
- If I continue the pregnancy and keep the child what will change in my life?
- Will I have support from my partner?
- If not, can I parent this child on my own?
- How do I feel about an ongoing connection with the conception partner, and possibly sharing custody of a child with this person if they are not my partner?
- Will I have practical support from family or friends?

- What effect will continuing the pregnancy and parenting have on my current children?
- Do I have secure housing? If not could my family help with this?
- Am I in a financial position to support myself and a child? Do I have significant debts, or could I manage ok?
- How will continuing the pregnancy affect my education?
- How will continuing the pregnancy affect my career and what I want in my career?
- What are my hopes and goals for my life?
- What would I like my life to be like in 1 year... 5 years...10 years.
- How will the decision I make about this pregnancy affect what I want for myself in the future?
- What are my main fears about continuing the pregnancy?
- What are my fears about ending the pregnancy?
- What strategies can I use to deal with those fears?

It's a good idea to take some time to sit down and focus on the questions that are relevant to you, and to write down your responses.

### What to do next if you are continuing the pregnancy

Make an appointment to see your GP. They will be able to organise the routine blood tests done in pregnancy, talk to you about pregnancy health, pregnancy care, and your birthing options. You will also need a referral from them for pregnancy care.

For more information about pregnancy care and birth options for women living in the ACT region see:

**'Having a Baby in Canberra'**  
[havingababyincanberra.org.au](http://havingababyincanberra.org.au)



## Adoption

Adoption is where the legal rights and responsibility for a child are permanently transferred from the birth parents to the child's adoptive parents. This means that once the child is adopted, the birth parents will no longer be the child's legal parents and the child's adoptive parents will become the child's legal parents. This is a permanent arrangement.

Adoption arrangements take place after the child is born; however, adoption services provide information and support for birth parents throughout the pregnancy. In the ACT Child and Youth Protection Services, within the Community Services Directorate, is responsible for providing adoption services in accordance with the Adoption Act 1993.

Both parents of a child have legal rights and ideally in most situations, both parents should be involved in the adoption plan for their child. Sometimes, however, it may not be possible for a mother to involve the child's father in the adoption decision.

An adoption decision for a child only becomes official when the parent signs a consent form. This happens after the child is born. The parent cannot sign a consent form until the baby is at least 8 days old and they must have had this consent form for at least 14 days. There is then a cooling-off period of 30 days during which the birth parent can change their mind.

When birth parents make an adoption plan for their child, they may choose to be actively involved in the process of selecting adoptive parents. Birth parents may choose to meet the adoptive parents and to have occasional contact with the child and adoptive parents. Photographs and updates about the child can also be provided to the birth parents by the adoptive parents. Any contact arrangements, if agreeable to both parties, can be included in the conditions of adoption order made in the Supreme Court.

Adoption can be a challenging process, and it is important that you get good support and counselling if you are considering this option.

### Questions you may want to ask yourself if you are considering adoption

- How do I feel about adoption?
- What do I feel the pros and cons of adoption are? (write this down)
- How do I feel about each of these pros and cons. (write these down too)
- How would I feel putting a child up for adoption?
- How do I think I may feel in a year from now if I put a child up for adoption?
- How do I think I may feel in 5 or 10 years from now if I put a child up for adoption?
- Do I know anyone who has put a child up for adoption?
- Are there options for temporary care of a child that I want to consider?
- What are my main fears and uncertainties about adoption?
- What strategies can I use to deal with those fears and uncertainties if I decide to place my child for adoption.

### What to do next if you want to place a child for adoption

Gather as much support as you can from people that you trust, and access counselling services as needed. Talk to the conception partner if that is possible and safe. Gather more information about adoption and talk to the people who work in placement of children for local adoptions.

There are options for temporary care (such as foster care or kinship care) which may be an alternative to adoption. You can discuss this with the staff who work with adoptions.

**Further information:** ACT Community Services Website: [communityservices.act.gov.au](http://communityservices.act.gov.au) and search for Adoptions. You can also contact Child and Youth Protection Services via phone on 02 6207 1466 or email [adoptions@act.gov.au](mailto:adoptions@act.gov.au)

## Abortion

Abortion when a pregnancy is terminated using medication or a surgical procedure. It is estimated that 1 in 4 women in Australia have at least one abortion in their lifetime. Most of these abortions occur in early pregnancy and over 92% occur in the first 14 weeks of pregnancy.

### Types of abortion:

#### Medication abortion

Medication abortion (also called medical abortion) is available in Australia for pregnancies under 9 weeks. This is a non-surgical method of ending a pregnancy using medication to induce a miscarriage.

Medication abortion is available through abortion clinics and through some GPs who have received specialised training in providing medication abortion. In the ACT region there are some GPs who can provide medication abortion free for ACT residents. Please call SHFPACT on 02 6247 3077 for details.

#### How does medication abortion work?

Medication abortion uses two different medications to end a pregnancy. The first medication, taken at the time of your visit to the clinic or GP, blocks the hormone progesterone (this is the hormone that maintains the pregnancy). The second medication is taken 24-48 hours later and causes the uterus to expel the pregnancy tissue.

The process of expelling the tissue usually takes about 4 hours but can take up to a few days. You will experience strong period-like pain, and bleeding will occur as part of the process. The bleeding is heavier than for a period and there may be blood clots. You may also experience nausea, vomiting, diarrhoea, dizziness, fever, shaking and headache after taking the second medication.

The bleeding should ease within the first week, but lighter bleeding can continue for up to a month afterwards.

Success rates for medication abortion are up to 98%. The complication rate, while very low, is slightly higher than for surgical abortion.

For more detailed information regarding medication abortion see



**Children by Choice:**  
[www.childrenbychoice.org.au/information-support/abortion/medication-abortion-2/](http://www.childrenbychoice.org.au/information-support/abortion/medication-abortion-2/)

#### Surgical abortion

Surgical abortion is a simple surgical procedure which is carried out as a day procedure. While surgical abortion can be done under local anaesthetic (which is injected into the cervix) or general anaesthetic, it is most often performed under twilight sedation, where you are asleep during the procedure. Twilight sedation has few health risks and is considered safer than a general anaesthetic.

#### What happens with a surgical abortion?

First you need to make an appointment with the abortion service. They will advise you about how long you will need to fast beforehand and what to bring to the clinic with you. You may spend from two hours up to half a day at the clinic. While procedures vary with individual clinics this will usually include admission, consultations with a nurse, with an anaesthetist, and with the doctor performing the abortion, and may include a blood test and an ultrasound.

The actual abortion procedure takes only 10 to 15 minutes to perform and is done by putting a small tube through the cervix and gently removing the pregnancy tissue from the uterus using suction.

Following the procedure, you will spend some time recovering before you go home. Most clinics require you to have someone to drive you home because you have had sedation.

Later gestation abortions (over 14 weeks gestation) may require a different procedure, sometimes involving two separate visits to

the clinic, one for the preparation of the cervix and one for the procedure itself. Later gestation abortions require a more complex procedure and can therefore be more expensive, with cost increasing with the length of the pregnancy.

Availability of later gestation abortion and the cost involved varies in different states and territories. For more information about this call SHFPACT on 02 62473077.

For more detailed information regarding the process of surgical abortion see

#### Children by Choice:



[www.childrenbychoice.org.  
au/information-support/  
abortion/surgical-abortion/](http://www.childrenbychoice.org.au/information-support/abortion/surgical-abortion/)

### How safe is abortion?

In Australia abortion is performed by highly trained medical professionals. Surgical abortion is performed in accredited health facilities and is one of the safest medical procedures available. The complication rate for early surgical abortions is very low at around 1%. This rate rises to around 5% after 15 weeks' gestation. The complication rate for medication abortion is between 2% and 5%.

If complications do occur the most common are retained products (where a small bit of pregnancy tissue remains in the uterus after the abortion-usually requiring a second procedure) or infection, which is dealt with by using antibiotics.

We know that legal abortion has a very low rate of complications. It is a common procedure in Australia that forms part of good reproductive health care.

Research has shown that there is no link between safe, legal abortion and future problems such as infertility, breast cancer, or long-term mental health problems.

### How you might feel following an abortion

Most people cope very well after having an abortion. The most common feeling that is expressed is relief.

How you feel afterwards will depend on many factors, including your individual circumstances, how you felt about the decision, your individual beliefs, and values, if you had good support and someone to talk to, and importantly if the decision was yours.

Studies show that most people who have abortions (around 98 per cent) feel that they made the right decision.

It is important to know that when a pregnancy ends, whether it is in abortion, miscarriage, or birth, the pregnancy hormone levels drop steeply around 3 to 4 days later. This can make some people feel sad, empty, and weepy. If this happens it is helpful to remember that it is just the effect of the hormone levels dropping and will pass in 24 hours or so.

It is normal to experience a mix of feelings after an abortion. This may include relief, sadness, loss, regret, guilt, happiness, or anger. There is no right or wrong way to feel. If you find you are experiencing some persistent difficult feelings talk to a trusted support person or to a pregnancy options counselling service.

The very small number of people who do experience significant emotional problems after an abortion tend to be those who were very unsure about their decision at the time of the abortion or were persuaded or coerced into having an abortion by someone else.

Some people are pressured by partners,

### Myths about abortion

There are many myths about abortion. These are often put forward by individuals or groups who are anti-choice and who run scare campaigns about abortion, or by people who are simply misinformed. As a result, there is a lot of biased and inaccurate information, particularly online.

family, or others to have an abortion. If someone wants to continue the pregnancy, a coerced abortion can result in emotional problems for them afterwards.

If you have had problems with anxiety or depression in the past and found the decision to have an abortion very difficult, then you may need more support during this time. This support may be from family or friends or through professional counselling.

### **Availability of abortion in Australia**

Abortion is legal in all states & territories in Australia. However, the availability of abortion services and the limits of how far pregnant you can be when you can have an abortion vary.

In the Australian Capital Territory (ACT) abortion is a legal, regulated health service and is covered under the Health Act 1993.

### **Abortion services in the ACT region:**

**MARIE STOPES AUSTRALIA** has a clinic in Canberra city which provides medication abortion up to 9 weeks gestation and surgical abortion up to 14 weeks gestation at time of publication. Abortion is free through Marie Stopes Canberra for ACT residents.

Call 1300 003 707 or book online via their website: [mariestopes.org.au](http://mariestopes.org.au)

**GENERAL PRACTITIONERS:** There are GPs who can provide medication abortion free of cost for ACT residents. Please call SHFPACT on 62473077 or see My Pregnancy Options ACT online for details.

Later abortion is available through other services and is dependent on gestation. Please call SHFPACT on 62473077 for details.

If you need further information or are having difficulty accessing an abortion, please contact SHFPACT on 02 62473077 and ask to speak to one of our pregnancy options counsellors.

### **Questions you may want to ask yourself if you are considering abortion**

- How do I feel about the issue of abortion generally?
- How do I feel about abortion for myself?

- What are the pros and cons of abortion for me (write these down).
- How do I think I may feel 6 months from now?
- How do I think I may feel 2 years from now?
- Do I know anyone who has had an abortion?
- What are my main fears and uncertainties about having an abortion?
- What strategies can I use to deal with those fears and uncertainties?

### **What to do next if you decide to have an abortion**

You can call the abortion services directly to make an appointment; you do not need a doctor's referral for this. See My Pregnancy Options Act for contact details.

If you would like further information before making a booking with an abortion service, make an appointment with SHFPACT's free Pregnancy Options Counselling Service

### **Pregnancy Options Counselling**

Counselling is really an umbrella term, and can include advice, information, support, education and/or therapy.

Counselling can offer a valuable and much-needed resource when deciding about a pregnancy. Some may not feel a need for it, while others can find it very helpful.

Counselling can be whatever you need it to be. If you feel you need support with decision making, feel you have already decided and just need to talk it through, or simply need someone to talk to, counselling can be helpful.

Some people may feel uncomfortable or unable to talk with friends or family about an unintended or unwanted pregnancy. In this situation counselling can offer an opportunity to talk about your individual situation in a supportive non-judgmental environment. Counsellors can also provide accurate information which may help you in making the decision which is right for you at this time in your life.

## A word of caution

It is important that the counselling you access is non-judgmental, non-directive, and refers for all-options. Some organisations offering pregnancy support or counselling may misrepresent their service when advertising or on their websites and may be anti-choice organisations.

Always check that any support or counselling service is a genuine all-options, non-directive service before you go there. One way of checking is to ask if they will give you information about abortion services and refer you if needed. An all-options service will always do this.

## Pregnancy Options Counselling in the ACT region

**Sexual Health and Family Planning ACT (SHFPACT)** provides professional, unbiased, non-directive, respectful, and confidential counselling for those who are deciding whether to continue a pregnancy, or who are experiencing an unintended pregnancy.

This is a free service, and the only free non-directive, face to face pregnancy options counselling service in the ACT. The counselling service is staffed by trained experienced counsellors.

SHFPACT pregnancy options counsellors are also able to refer you to external counsellors should you need more complex counselling.

If you don't feel you need counselling but would like information about your options, our pregnancy options counsellors can also help you.



To make an appointment at SHFPACT's clinic or Pregnancy Options Counselling Service call (02) 62473077 or follow the QR Code.

## Other important contacts

### Parenting advice/support



**Having a baby in Canberra**  
[havingababyincanberra.org.au](http://havingababyincanberra.org.au)



**Parentlink**  
13 34 27  
[parentlink.act.gov.au](http://parentlink.act.gov.au)

### Adoption



**Child and Youth Protection Services**  
02 6207 1466  
[adoptions@act.gov.au](mailto:adoptions@act.gov.au)  
[communityservices.act.gov.au](http://communityservices.act.gov.au)

### Abortion



**My Pregnancy Options ACT**  
<https://mypregnancyoptionsact.org.au/find-a-service/>



**Marie Stopes Australia**  
1300 003 707  
[mariestopes.org.au](http://mariestopes.org.au)

### References:

- Children by choice: [childrenbychoice.org.au](http://childrenbychoice.org.au)
- Marie Stopes International: [mariestopes.org.au](http://mariestopes.org.au)
- Better Health Channel: [betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Abortion\\_counselling\\_options](http://betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Abortion_counselling_options)
- [betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Abortion\\_emotional\\_issues](http://betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Abortion_emotional_issues)
- [betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Abortion\\_some\\_misconceptions](http://betterhealth.vic.gov.au/bhcv2/bhcarticles.nsf/pages/Abortion_some_misconceptions)
- Office for Children, Youth and Family Support: [dhcs.act.gov.au/ocys/services/adoptions](http://dhcs.act.gov.au/ocys/services/adoptions)
- University of Iowa Maternity Centre: [uihealthcare.com/depts/maternitycenter/pregnancy/emotions](http://uihealthcare.com/depts/maternitycenter/pregnancy/emotions)
- The Royal Women's Hospital Victoria: [thewomens.org.au/Abortionthemylths](http://thewomens.org.au/Abortionthemylths)

SHFPACT is a health promotion charity (HPC) assisted with funding from the ACT Government. All donations to SHFPACT over \$2.00 are tax deductible.

Supported by

