

IT'S EASY TO GET A

DISCOUNTED CONSULTATION

WE ARE MAKING OUR SERVICES MORE AFFORDABLE FOR EVERYONE

BECOME A MEMBER AND SAVE TODAY!

WHAT DO YOU GET AS A MEMBER?

- Special members discount rates for consultations at SHFPACT clinic and Canberra Menopause Centre.
- 5% discount on bookshop purchases.
- Free or discounted attendance at health information and special events. For example: Healthy Menopause Information Evenings by the Canberra Menopause Centre, Parent Information Evenings about communicating with your children and young people about puberty, relationships, etc.
- SHFPACT members enews that will keep you updated with special members events, activities and other services.

JOIN ONLINE AT shfpact.org.au OR AT RECEPTION



OPENING HOURS

Please contact our reception staff on 02 6247 3077 or visit shfpact.org.au for SHFPACT clinic hours

LOCATION

SHFPACT is conveniently located in the city.
Level 1,
28 University Avenue
Canberra ACT 2601

CONTACT

02 6247 3077
shfpact@shfpact.org.au
0400 770 999 (SMS ONLY)

ENGAGE

shfpact.org.au
facebook.com/shfpact
twitter.com/shfpact

DONATE

All donations to SHFPACT over \$2.00 are tax deductible

SHFPACT CLINICAL SERVICES

- Contraceptive advice, information and services
- IUDs and other long acting contraceptive options
- Cervical screening
- Canberra Menopause Centre
- Unplanned Pregnancy Counselling Service
- Sexually transmissible infections (STIs): testing, treatment and advice
- Blood borne virus and HIV testing
- Emergency contraception (the 'Morning After' Pill)
- Breast checks and breast awareness education
- Fertility issues and pre-conception advice
- Sexual difficulties and sexuality education
- PrEP information, advice, and prescriptions

SHFPACT is a health promotion charity (HPC) assisted with funding from the ACT Government.

I'M ALL ABOUT THE PILL



Sexual Health and Family Planning ACT
Level 1, 28 University Ave, Canberra City
Call 02 6247 3077 • SMS 0400 770 999
or email shfpact@shfpact.org.au



shfpact.org.au

Last updated February 2020

WHAT IS THE COMBINED ORAL CONTRACEPTIVE PILL?

The combined oral contraceptive pill, usually simply called 'the pill', is a contraceptive pill taken by mouth which contains two hormones, oestrogen and a progestogen.

HOW DOES IT WORK?

The pill works by preventing ovulation, which means it stops the ovaries from releasing an egg each month, which in turn means that fertilization cannot occur and a pregnancy cannot begin.

HOW EFFECTIVE IS IT?

The pill is approximately 93% effective.

HOW IS IT TAKEN?

The pill is taken by mouth every day. Most pill types come in a box of three to four sachets with each sachet containing a month's supply of pills. Each pill sachet contains hormone pills that you usually take for three weeks, and inactive or sugar pills (which help you to keep the habit of taking the pill every day) that you take for seven days before starting the hormone pills again. You can also take the hormone pills for longer than three weeks so that you skip periods. If you think you would like to do this discuss it with your doctor.

WHO CAN TAKE THE PILL?

The pill is suitable for most contraceptive users.

There are some medical conditions which make the pill unsuitable, these include the following:

- Deep venous thrombosis (blood clot), stroke or heart attack.
- Severe liver problems.
- Migraine with aura (visual or other disturbances starting just before the migraine).
- Some auto immune conditions that increase the risk of blood clots.

The pill may also be unsuitable:

- If you have high blood pressure, diabetes, gall bladder disease, active liver disease, kidney disease, some blood problems, or if you are on some other medications which may interact with the pill.
- If you have a history of breast cancer or cervical cancer.
- If you are over 35 and smoke.
- If you have unexplained bleeding from the vagina.

If any of these conditions apply to you, let your doctor help you decide if the pill is suitable for you.

ADVANTAGES OF THE PILL?

- It is very effective with correct use.
- It is readily accessible.
- It is easily reversible.
- It gives you a predictable bleeding pattern and allows you to skip periods if you want.
- It can be useful in controlling heavy or painful periods and managing Premenstrual Syndrome.
- It can improve acne.

DISADVANTAGES OF THE PILL?

- It must be taken every day.
- It is less effective if not used perfectly (for example if pills are missed).
- Some pill types can be expensive.
- Effectiveness may be reduced by vomiting or diarrhoea.

THE PILL AND YOUR HEALTH

Serious health problems caused by the pill are very rare — the most significant of these is blood clotting. Symptoms of this are severe sudden chest pain, severe pain or swelling in one leg, sudden blurred vision or loss of sight, slurring of speech or sudden severe headache. If you have any of these symptoms contact your doctor immediately.

Your risk of developing a blood clot is increased when you are not mobile for an extended period of time e.g. sitting in a car or airplane on a long trip, or if you are planning to have surgery. Please talk to your doctor if you are planning any of these.

As well as being a contraceptive, the pill has other health benefits. When on the pill you are less likely to develop a serious pelvic infection, cancer of the ovary, cancer of the endometrium (lining of the womb), cancer of the bowel, anaemia, non-cancerous breast lumps, and cysts of the ovary.

When you are taking the pill, periods are generally less painful, the bleeding is usually lighter and more regular, and there is often less premenstrual tension. Acne may also improve.

Most people feel fine on the pill, but it is common to have some minor side effects initially. You might get some bleeding in between periods, sore breasts and mild nausea (feeling sick) for the first couple of months. These side effects usually settle by themselves.

Some people report weight changes, reduced desire for sex, and mood changes, however, there is no scientific evidence to show that the pill causes these effects, other aspects of life may be contributing factors.

Very rarely the oestrogen in the pill can cause patchy brown discolouration of the skin on the face called melasma. This is more noticeable if you spend a lot of time in the sun. If melasma does occur it usually does so in the first few months after starting the pill. If you notice this discolouration starting to appear on your face see your doctor.

TYPES OF PILLS AVAILABLE

There are quite a few different types of the pill available. There are some differences between them and it may take a few trials to find the one most suitable for you. Don't be afraid to discuss the different options with your doctor.

STARTING THE PILL

As long as you are absolutely certain that you are not pregnant, you can start the pill at any time of your menstrual cycle. If you start with a hormone pill on Day 1 to Day 5 of your cycle (Day 1 is your first day of bleeding), you are immediately protected from pregnancy.

If you start on any other day of your menstrual cycle, you will need to use additional contraception, such as condoms, or abstain from intercourse, until you have taken 7 of the hormone pills.

WHAT TO DO IF YOU MISS A PILL

Is it more than 24 hours since the pill should have been taken? (This means more than 48 hours since you last took a pill).

